

Claims

1. Macromolecular photocrosslinkers having a general formula

3-16-00

7/1

C-m P units

5 (A)_n(B)_m(C)_p, wherein

(i) A, B and C are units of substituted ethylene or siloxane groups in the macromolecular structure;

(ii) C carries a photoactive groups;

10 (iii) n = 0-98 mole %, m = 0-98 mole %, n+m = 50-98 mole % and p = 0.5-50 mole %;

and when said photoactive groups are exposed to light of determined wavelengths above 305 nm, radicals are generated and retained on the macromolecular photocrosslinkers and reacting so as to accomplish a crosslinked network structure.

cage effect
Angewandte
chemie

15

2. Photocrosslinkers according to claim 1 characterized in that said photoactive group comprises a phosphine oxide.

3. Photocrosslinkers according to claim 2 characterized in that the photoactive group is an acyl- or aroyl phosphine oxide.

20

4. Photocrosslinkers according to claim 3 characterized in that the photoactive group is linked to the ethylene groups of units C by a linking group comprising a phenylene group, said phenylene group being optionally substituted.

25

5. Photocrosslinkers according to claim 1, wherein the ethylene units A, B, C of the macromolecular structure comprises substituents in accordance with:

A = -CH₂-C(R¹R²)-, B = -CH₂-C(R¹R³)-, C = -CH₂-C(R¹R⁴)-, wherein

30

AMENDED SHEET

17/11

09036647-025502

WO 00/55212

PCT/EP00/02539

26

R^1 is hydrogen or methyl;

R^2 is $-\text{CON}(\text{Me})_2$, $-\text{CO}_2\text{CH}_2\text{CH}_2\text{OH}$, $-\text{OCOCH}_3$, $-\text{OCOCH}_2\text{CH}_2\text{Ph}$, $-\text{OH}$ or a lactam group;

R^3 is $-\text{CON}(\text{Me})_2$, $-\text{CO}_2\text{CH}_2\text{CH}_2\text{OH}$, $-\text{OCOCH}_3$, $-\text{OCOCH}_2\text{CH}_2\text{Ph}$, $-\text{OH}$ or a lactam group when B is $-\text{CH}_2-\text{C}(\text{R}^1\text{R}^3)-$ with the proviso that R^2 and R^3 are not the same unless R^2 and R^3 is $-\text{OH}$; and

R^4 is $-\text{R}^5\text{C}(\text{O})\text{P}(\text{O})\text{R}^6\text{R}^7$ or $-\text{R}^5\text{P}(\text{O})\text{R}^6\text{OC}(\text{O})\text{R}^7$, wherein R^5 , R^6 and R^7 are selected among same or different aryl groups comprising phenyl, methylphenyl, dimethylphenyl, trimethylphenyl, methoxyphenyl, dimethoxyphenyl, trimethoxyphenyl, methylolphenyl, dimethylolphenyl, trimethylolphenyl or styryl radicals.

6. Photocrosslinkers according to claim 5, wherein R^2 and R^3 are selected so as to form a water-soluble molecule.

7. Photocrosslinkers according to claim 5, wherein said lactam units together with units A or B constitute N-vinylpyrrolidone units.

8. Photocrosslinkers according to claim 5, wherein at least one of R^2 and R^3 is hydroxyl.

9. Photocrosslinkers according to claim 5, wherein A is N-vinylpyrrolidone, B is vinyl alcohol.

10. Photocrosslinkers according to claim 1 ~~or 5~~ provided with functional groups for crosslinking.

11. Photocrosslinkers according to claim 10 provided with functional groups selected among vinylic, acrylic and methacrylic groups.

12. Photocrosslinkers according to claim 1 characterized in that units A, B and C are

Amend
A1

WO 00/55212

PCT/EP00/02539

27

siloxane monomer units of a general formula $-R_aR_bSiO-$, wherein R_a and R_b in units A and B are selected among lower substituted or unsubstituted alkyl groups, aryl groups and arylalkyl groups.

13. Photocrosslinkers according to claim 12, wherein at least one of R_a and R_b is an aryl or arylalkyl group.

14. Photocrosslinkers according to claim 13, wherein at least one of R_a and R_b is substituted with one or more fluorine atoms.

15. Photocrosslinkers according to claim 1, wherein units A, B, C are siloxane units comprising substituents in accordance with:

A is $-Si(R^1R^2)-O-$, B is $-Si(R^1R^3)-O-$ and C is $-Si(R^1R^4)-O-$, wherein

R^1 is C1 to C6 alkyl; R^2 is C1 to C6 alkyl or phenyl; R^3 is R^1 , R^2 or C1 to C6 fluoroalkyl;

→ R^4 is $-R^5R^6C(O)P(O)R^7R^8$ or $-R^5R^6P(O)R^7OC(O)R^8$, wherein R^5 is a spacing group; R^6 , R^7 and R^8 are selected among same or different aryl groups comprising phenyl, methylphenyl, dimethylphenyl, trimethylphenyl, methoxyphenyl, dimethoxyphenyl, trimethoxyphenyl, methylolphenyl, dimethylolphenyl, trimethylolphenyl or styryl radicals.

16. Photocrosslinkers according to claim 15, wherein R^5 is aliphatic spacing group comprising between one and ten atoms.

17. Photocrosslinker according to claim 16, wherein said spacing group is $(-CH_2)_n$, wherein n is between 1 and 10.

18. Photocrosslinkers according to claim 15, wherein R^1 is methyl; R^2 is methyl or phenyl; R^3 is R^1 , R^2 or $-CH_2CH_2CF_3$.

WO 00/55212

PCT/EP00/02539

28

19. Photocrosslinkers according to claim 15 having functional acrylic groups in its terminal ends.

Amend 20
A2
50.5/50+
20. A method of forming a macromolecular crosslinked network from a composition comprising a photocrosslinker according to any of claims 1 to 19 by irradiating said composition with light exceeding a wavelength of about 305 nm for a time sufficient to form a solid article.

21. A method forming a macromolecular crosslinked network from a composition comprising a photocrosslinker according to any of claims 1 to 11 and at least one copolymerizable vinyl, acrylic or methacrylic monomer.

22. A method according to claim 20, wherein said composition further comprises a polymer provided with functional vinyl, acrylic or methacrylic groups.

23. A method according to claim 22, wherein said polymer has a backbone of ethylene units.

24. A method according to claim 22, wherein said polymer is a polysiloxane. 1712

Amend
A3
25. A method according to any of claims 20 to 24, wherein an ophthalmic lens is produced.

112 no ante
128 ?
26. A method according to claim 25, wherein an intraocular lens is produced in the capsular bag of the eye.

Amend
A4
27. An ophthalmically acceptable composition comprising photocrosslinkers according to any of claims 1 to 19, having a refractive index greater than about 1.39 and a viscosity such that said composition can be injected through standard cannula having a needle of 15 Gauge, or finer.

WO 00/55212

PCT/EP00/02539

29

ay

28. The use of ^{see *And*} photocrosslinkers according to any of claims 1 to 19 in an
ophthalmologically acceptable composition for injection into the capsular bag of the eye.

205220" 4993660